

ELECTRIC SHOCKS ARE EFFECTIVE IN TREATMENT

OF LETHAL EFFECTS OF RATTLESNAKE ENVENOMATION
IN MICE

EDWARD K. JOHNSON,¹ KENNETH V. KARDONG^{2*} and STEPHEN P. MACKESSY¹

¹Department of Physiological Sciences, Oklahoma State University, Stillwater, Oklahoma 74078 U.S.A., and

60 sec of the venom injection. None of the shocked experimental mice survived at greater

frequency than the control group. Twenty four hour survival of envenomated control and shocked mice were as follows: at 6.0 mg/kg control (4/4) shock (4/4); at 3.0 mg/kg

control (4/4), shock (4/4); at 1.75 mg/kg, control (6/8), shock (7/8); at 1.5 mg/kg, control (0/4), shock (0/4). Parentheses indicate number dead/total number tested, respectively. Thus we found no evidence that high voltage, low ampage electric shock is

efficacious in treating the lethal effects of rattlesnake envenomation in mice. There are two possible reasons for this apparent absence of effect in mice.

- PARRISH, H. M. (1959) Poisonous snakebites resulting in lack of venom poisoning. *Va. med. Mon.* **86**, 396.
- REID, H. A. (1982) Animal poisons. In: *Manson's Tropical Diseases*, 18th Edn (MANSON-BAHR, P. E. C., Apted, F.I.C., Eds). London: Bailliere Tindall.