toxins

Review

Colubrid Venom Composition: An -Omics Perspective

Inácio L. M. Junqueira-de-Azevedo [1,*], Pollyanna F. Campos [1], Ana T. C. Ching [2] and Stephen P. Mackessy [3]

1 Laboratório Especial de Toxinologia Aplicada, Center of Toxins, Immune-Response and Cell Signaling (CeTICS), Instituto Butantan, São Paulo 05503-900, Brazil; pollyanna.fc@gmail.com
2 Laboratório de Imunoquímica, Instituto Butantan, São Paulo 05503-900, Brazil; anatung@gmail.com
3 School of Biological Sciences, University of Northern Colorado, Greeley, CO 80639-0017, USA; stephen.mackessy@unco.edu
* Correspondence: inacio.azevedo@butantan.gov.br; Tel.: +55-11-2627-9731

Academic Editor: Bryan Fry
Received: 7 June 2016; Accepted: 8 July 2016; Published: 23 July 2016

Abstract: Snake venoms have been subjected to increasingly sensitive analyses for well over 100 years, but most research has been restricted to front-fanged snakes, which actually represent a relatively small proportion of extant species of advanced snakes. Because rear-fanged snakes are a diverse and distinct radiation of the advanced snakes, understanding venom composition among “colubrids” is critical to understanding the evolution of venom among snakes. Here we review the state of fact present in several colubrid venoms, while others are only transcribed, at lower levels. This work provides insights into the evolution of these toxin classes, but because only a small number of species have been explored, generalizations are still rather limited. It is likely that new venom protein families await discovery, particularly among those species with highly specialized diets.

Keywords: Colubridae; evolution; proteins; snake; toxins; transcriptomics

1. Introduction

More than one hundred years of biochemical and pharmacological studies have resulted in an exceptional depth of knowledge about snake venoms. The major toxins of the most medically important taxa of venomous snakes were determined by rst generation approaches including protein chemistry, comparative pharmacology and cladistics methods borrowed from evolutionary biology. Advances in molecular biology, particularly protein and nucleic acid sequencing techniques, greatly

Table 2. Minor snake venom components and their occurrences in colubrid species.

Species	Enzymatic							Non-Enzymatic										Reference
	5NUCL	AChE	DPP	FactV	FactX	HYAL	PDE	AVIT	bPLA2i	CVF	CYST	gPLA2i	KU-WA	NGF *	OHA	VEGF-A **	WAP	
Boiga dendrophila																		[42,43]
Boiga irregularis		T				t	t		t	t	t	t				t	t	[12,44,45]
Borikenophis portoricensis		B					B											[46,47]
Cerberus rynchops																		[48]
Coelognathus radiatus																		[49]
Dispholidus typus																		[50–52]
Erythrolamprus miliaris		t	t			t				t		T				t	xt	This work; [50]
Erythrolamprus poecilogyrus																	x	[50,51]
Helicops angulatus																		[53]
Hypsiglena sp.			t										tP		t	t	t	[13]
Hypsiglena torquata																		[54]
Leoiheterodon madagascarensis																		[50]
Macropisthodon rudis							t											[55]
Opheodrys aestivus		t	t							t	x		t	x	t	t	t	[9]
Oxybelis fulgidus																		[56]
Oxyrhopus guibei		t	t							t								

Table 3. Putative new snake venom components identified from colubrid species.

Species	Enzymatic		Non-Enzymatic	Reference
	svLIPA	PLA		

Figure 1. Schematic cladograms showing the phylogenetic relationships among families and species of snakes discussed in this work (colored branches). The cladogram was based on the phylogenetic tree proposed by Pyron et al. [34]. Dashed lines in Philodryas indicate the presumed placement of P. chamissonis

It is interesting to note that most -omics characterizations of colubrid venoms have addressed

In colubrids, they seem to not be among major components and have been detected in only a few species. In the colubrine Trimorphodon biscutatus, an enzyme was puri ed and its partial sequence indicated that it was a type IA PLA $_2$ [61]. However, in another colubrine (Dispholidus typus) [51], in the dipsadine Oxyrhopus guibei (this work) and in the pseudoxyrhophiine (family Lamprophiidae) Leioheterodon madagascariensis [51], among others, the reported type of PLA $_2$ is IIE.

The occurrence of transcripts coding for enzymes of IIE subtype in the venom glands indicates a possible independent recruitment of a PLA $_2$ to the venom, since they are distinct from the type IIA paralogs commonly expressed in the venom glands of viperid snakes [51]. Whether or not these type IIE PLA $_2$s represent truly new toxins or accessory proteins of the venom glands remains to be clari ed, but Hargreaves et al. [9] found them to be exclusively expressed at low levels in the venom glands of the species tested.

Despite being very common in the venom of other groups of snakes, L-Amino Acid Oxidase (LAAO) was thought to be essentially non-existent in colubrid venoms (e.g., [71]). Very low levels of LAAO activity were detected in Brown Treesnake (B. irregularis) venom [72]; however, assays of venom from 13 different species of colubrine, dipsadine and natricine rear-fanged snakes detected no LAAO activity [73]. In a comparative transcriptomic analysis of tissues from Pantherophis guttatus, an LAAO was shown to be expressed in the scent gland but not in the salivary glands of this species [9], suggesting it is not a venom component. Recently, however, an LAAO was found moderately expressed in the venom glands of the colubrid Phalotris mertensi, and the MS/MS spectrometric analysis clearly showed its presence in the venom of this species [57].

2.3. Major Snake Venom Non-Enzymatic Components

Three-Finger Toxins (3FTx) are major constituents of Elapidae venoms and represent the lethal component of the majority of species of this family. These toxins seem to have differential importance in different subfamilies of colubrids. Alpha-colubritoxin from Coelognathus radiatus was the rst colubrid toxin isolated and sequenced [49] and several other 3FTx, such as denmotoxin and irditoxin, functionally characterized in members of the subfamily Colubrinae, were demonstrated to be abundant toxins with taxon-speci c activities [42,45]. The -omics characterization of Boiga irregularis venom showed that 3FTx dominate the transcriptome of this species (67.5% of toxin transcripts) [12]. The authors described 58 unique 3FTx sequences grouped into at least 10 sequence clusters that were proteomically con rmed in the venom. These clusters could be arranged in three groups based on the structural characteristics, but none of them were closely related to the above-mentioned irditoxin from the same species. Together with individual sequences isolated from other genera [9,50,56], 3FTx seem to be major components in many venoms of the subfamily Colubrinae. In contrast, in the Dipsadinae, 3FTx are not found [59,60] or are detected at minor abundance and diversity levels [57]. However, in the current work, we retrieved sequences from the transcriptome of the Duvernoy's venom gland of Xenodon merremi (a dipsadine colubrid) that were expressed at high level (Table 1). 3FTx-like sequences were also reported in venom glands of the family Lamprophiidae, as well as in species at the base of the Alethinophidia snake radiation (including species in the Cylindrophiidae and Pythonidae [50]). Nevertheless, it appears likely that 3FTx-like transcripts found in gland tissue of these latter two families may represent house-keeping genes, rather than toxins [9,74].

C-Type Lectins (CTL) are ubiquitous venom components in many snake groups, and they are also found abundantly in colubrid venoms. In Colubridae, amounts of venom CTL transcripts vary from 2% in Philodryas olfersi [59] to as much as 21% in Phalotris mertensi of total transcripts [57]. They were also reported in the snakes Pseudoferania polylepis and Cerberus rynchops (family Homalopsidae) and were highly expressed in Cerberus [48]. From a phylogenetic tree of all colubrid CTLs and related orthologs (Figure 3), it is possible to observe the existence of distinct types of CTLs in non-front fanged snakes, although the phylogenetic reconstruction failed to resolve the evolutionary relationships among them. Nevertheless, in addition to the presence of a CTL-like (snalec) clade, largely found in colubrids, colubrid sequences are observed to be nested within the clade of Elapidae and Viperidae “true” CTLs

sequences (i.e., those with a predicted galactose binding motif QPD substituting the plesiotypic motif EPN: [75,76]. One of them (PMERREF_CTL04) was confirmed in *P. mertensi* venom and in fact has the QPD motif, indicating that predicted galactose-binding lectins should also be present in other Colubridae venoms. Moreover, some orphan transcripts observed in the venom glands of snakes from different families clustered completely outside of the clades of typical venom CTLs (Figure 3). Some of them are suggested to code for venom proteins, such as two transcripts highly expressed and proteomically detected in the venom of *P. mertensi* and similar transcripts are expressed at moderate levels in the venom glands of other colubrids (*Hypsiglena* sp. and *Boiga irregularis*; [12]), viperids (*Bothrops insularis*; [77]) and elapids (*Hoplocephalus bungaroides*; [78]). Interestingly, the encoded proteins from the transcripts of this group present not a single but various motifs (QPD, EPD, EPN, RPS, QVE, and EPK) for sugar binding at the second loop of the carbohydrate recognition domain. It indicates that these genes may have undergone a diversification process that parallels that experienced by other CTL types, i.e., the substitution of the binding motif of the original sugar ligand, mannose, by binding motifs to other types of carbohydrates.

Figure 3. Maximum likelihood circular cladogram showing the relationship among representative CTLs from different snake families. Bootstrap values are plotted close to internal nodes. Colors at the terminal nodes (circles) indicate typical vs. atypical venom proteins and the evidence of occurrence in the venoms. Colors in the diagram surrounding the cladogram indicate the taxonomic groups. The carbohydrate binding motifs, as discussed in the text (EPN, QPD, etc.), are indicated by red type. The protein sequences are referred by their GenBank accession numbers, except those initiated by the codes EMILISO, OGUIISO, PMERREF and XMERCLU, which are mentioned in the "definition" field of sequence files deposited in TSA.

and Phalotris mertensi [57]. In the latter, the corresponding protein was detected by shotgun MS/MS analysis of the venom, suggesting it may be a valid colubrid venom component. The colubrid proteins have a highly conserved signal peptide, almost identical to that of crotamine (see Supporting Figure 3 from [60]); the mature polypeptides display the same cysteines involved in the disulfide arrangement of crotamine, but the other residues are highly variable, making it difficult to establish the evolutionary relationship between them.

Kunitz-type proteins appear in snake venoms in several forms, sometimes as single-product precursors (KUN-1), at other times with tandem repeated domains (KUN-2), and less frequently associated with WAP domains in a protein designated ku-wap-fusin (KU-WA-FU) [87]. Although in some species of colubrids these components have a transcriptional level not indicative of a relevant participant in venom, in at least two species, Hypsiglena sp. [12] and Phalotris mertensi [57], they have medium or elevated expression levels and were also detected in the venom. In

Cobra venom factor (CVF) was clearly demonstrated as a venom component only in elapid snakes [93]. Although very similar sequences could be found expressed at low levels in some colubrids, the absence of protein detection in their venom suggests the transcripts could also be the endophysiological complement factor C3 expressed by blood cells within the venom glands.

Enzymatic inhibitors that typically function to protect snakes from the bites of other snakes are mainly produced in the liver and secreted into the plasma of venomous and non-venomous snakes [94], but some of them seem to be produced in the venom glands. For example, a specific paralog of a gamma-PLA_2 inhibitor ($gPLA_2i$) was shown to be exclusively expressed in venom glands of B. jararaca (Viperidae) [10

Figure 5. Maximum likelihood tree showing the relationship among svMMPs from different snake families and MMPs from other vertebrate groups. Bootstrap values are plotted close to internal nodes. The domain arrangement of each precursor type is depicted on the right. The types of evidence for the occurrence in venoms are indicated by “T” (transcribed) and “V” (detected in venom). The protein sequences are labeled by their accession numbers in GenBank, except those initiated by the codes EMILREF, OGUIISO, and PMERREF, which are mentioned in the “de nition” eld of sequence les deposited in TSA.

Figure 6. Maximum likelihood tree showing the relationship among lipocalin proteins from different snake families and from other vertebrate groups. Note that transcripts highly expressed in venom glands are all in the same clade. Bootstrap values are plotted close to internal nodes. The protein sequences are labeled by their accession numbers in GenBank, except those initiated by the code EMIL, which is mentioned in the "de nition" eld of sequence les deposited in TSA and the sequence OGUIREF_Lipo1 (Accession Number KX450875). Sequences labeled GAMF from *A aterrima* were translated from the original nucleotide contigs retrieved from TSA.

3. Conclusions

It is now abundantly clear that the venoms produced among the colubrid rear-fanged snakes are homologous with the much better characterized venoms of the front-fanged snakes. As trophic adaptations that facilitate feeding, venoms vary in composition with several important factors, including phylogeny, and so it is to be expected that among the diverse colubrid lineages, novel compounds, and new functional variants of better-known venom proteins, will be encountered. Much progress toward understanding rear-fanged snake venom composition has been made in the last decade, but, as indicated above, we have barely begun to explore the diversity of advanced snakes that comprise the colubrids. Transcriptomic and genomic approaches will greatly facilitate this work, but it must be remembered that functional assays should also accompany analysis of any venom, because the common recurring motif in venom biochemistry is to make the most of a stable molecular scaffold, perhaps best exempli ed by the varied pharmacologies of the three- nger toxin superfamily. These small, structurally conservative peptides have very similar crystal structures but affect systems as diverse as neurotransmission, the blood clot cascade, ion channel function, and salamander limb

for", and, to nd truly novel toxins that will likely be present in some colubrid venoms, we will have to look beyond the "normal" families of venom proteins.

4. Materials and Methods

4.1. Original Transcriptomic Data

4.1.1. Animals

Three specimens of Erythrolamprus miliaris (one male and two females) ve specimens of

4.1.3. Expressed Sequence Tags (ESTs) Generation

Xenodon merremtranscriptome was investigated by means of EST generation, prior to the common use of NGS (next generation sequencing). The pairs of glands from Xenodon merremi specimens were ground into a powder in liquid nitrogen and homogenized using a Polytron Tissue Homogenizer (Kinematica, Luzern, Switzerland). Total RNA was extracted with TRIZOL reagent (Life Technologies, Thermo Fisher Scienti c, Carlsbad, CA, USA) and mRNA was prepared using a column of oligo-dT cellulose (GE). To obtain 5

Con icts of Interest: The authors declare no con ict of interest.

Abbreviations

3FTx	three nger toxin
5NUCLEO	5′-nucleotidase
AChE	acetylcholinesterase
AVIT	AVIT protein
bPLA2i	beta type phospholipase A_2

7. Chapeaurouge, A.; Reza, M.A.; Mackessy, S.P.; Carvalho, P.C.; Valente, R.H.; Teixeira-Ferreira, A.; Perales, J.;

49. Fry, B.G.; Lumsden, N.G.; Wüster, W.; Wickramaratna, J.C.; Hodgson, W.C.; Kini, R.M. Isolation of a neurotoxin (alpha-colubritoxin) from a nonvenomous colubrid: Evidence for early origin of venom in snakes. J. Mol. Evol. 2003, 57, 446–452. [CrossRef] [PubMed]
50. Fry, B.G.; Scheib, H.; van der Weerd, L.; Young, B.; McNaughtan, J.; Ramjan, S.F.; Vidal, N.; Poelmann, R.E.; Norman, J.A. Evolution of an arsenal: Structural and functional diversi cation of the venom system in the advanced snakes (Caenophidia). Mol. Cell. Proteom. 2008, 7, 215–246. [CrossRef] [PubMed]
51. Fry, B.G.; Scheib, H.; Junqueira-de-Azevedo, I.L.M.; Silva, D.A.; Casewell, N.R. Novel transcripts in the maxillary venom glands of advanced snakes. Toxicon 2012, 59, 696–708. [CrossRef] [PubMed]
59

87. Doley, R.; Pahari, S.; Reza, M.A.; Mackessy, S.P.; Kini, K.M. The gene structure and evolution of ku-wap-fusin (Kunitz Waprin Fusion Protein), a novel evolutionary intermediate of the Kunitz serine protease inhibitors and waprins from Sistrurus catenatus