Venom phenotypes of the Rock Rattlesnake (Crotalus lepidus) and the Ridge-nosed Rattlesnake (Crotalus willardi) from M exico and the United States

Anthony J. Saviola [a,1], Anthony J. Gandara [a,1], Robert W. Bryson Jr. [b], Stephen P. Mackessy [a,*]

[a] School of Biological Sciences, University of Northern Colorado, Greeley, CO, USA
[b] Department of Biology, Burke Museum of Natural History and Culture, University of Washington, Seattle, WA, USA

a r t i c l e i n f o

Article history:
Received 24 May 2017
Received in revised form
21 July 2017
Accepted 17 August 2017
Available online 19 August 2017

Keywords:
Enzyme
Evolution
Phenotypic variation
Mexican highlands
Proteome
Toxin

2.3. Enzyme assays

Venoms from each of the four C. lepidus subspecies and four C. willardi subspecies were assayed for activity of enzymes commonly found in rattlesnake venoms (Mackessy, 2008, 2010a). Metalloproteinase activity was assessed using azocasein as a sub-



Fig. 1. Comparison of SDS-PAGE patterns of four subspecies of Crotalus lepidus (A) and Crotalus willardi (B) on NuPage gels (20 μg/lane). Typical protein families of bands of specific masses are indicated on the right; mass standards (MW Stds) are given in kilodaltons. Abbreviations: CTL – C-type lectin; LAAO – L-amino acid oxidase; PDE – phosphodiesterase; PLA_2 – phospholipase A_2; SVMP – snake venom metalloproteinase (PI and PIII classes); SVSP – snake venom serine proteinase. Note that PI-SVMPs are extremely faint to completely absent in C. l. klauberi and C. l. lepidus samples.

latter assay, the only comparison that was signi cantly different was between C. w. amabilis and C. l. klauberi ($p < 0.05$), whereas all other subspecies comparisons were not signi cant. Results of the

venom was also the most toxic toward lizards (0.17 mg/g) and the least toxic toward crickets (4.70 mg/g). Venoms from C. l. morulus showed the opposite pattern, being the least toxic toward lizards (6.68 mg/g) and the most toxic towards crickets (2.25 mg/g). Among the four C. willardi subspecies, venoms from C. w. obscuruswere more toxic toward mice, with an LD_{50} value of 1.58 mg/g. Of the four subspecies, C. w. willardi venom was the most toxic toward lizards (1.13 mg/g) while the venom from C. w. silus was the most toxic toward crickets (1.94 mg/g).

Venoms from C. w. amabilis showed lower toxicity toward crickets (10.48 mg/g) and lizards (6.65 mg/g). The numerical results of toxicity assays are summarized in Table 2.

3.4. Fibrinogenolytic activity

Venoms from C. l morulus and C. l. maculosusshowed potent brinogenolytic activity and completely degraded the alpha and beta brinogen subunits within 1 min of incubation (Fig. 4A). Venoms from C. l. klauberi and C. l. lepidusalso degraded the alpha and beta subunits of brinogen, after 5 min incubation (Fig. 4A). Similarly, venoms from C. w. willardi completely digested the alpha and beta subunits within the rst 5 min of incubation, whereas venoms from C. w. silusdigested the alpha subunit within the 5 min of incubation, but the beta subunit, although appearing to show a noticeably lower concentration after 30 min incubation, was not completely degraded even after 60 min (Fig. 4B). The degradation pattern of C. w. amabilis was similar to that observed for C. w. silus, with the alpha subunit being completely degraded after 1 min of incubation, while the beta subunit was not completely degraded until 30 min. The brinogenolytic pattern produced by venom of C. w. obscuruswas clearly different than those of the other three subspecies, and the alpha subunit of brinogen was not completely degraded until after 10 min of incubation. The beta subunit was stable for 30 min and began to show signs of degradation only after 60 min of incubation. None of the venoms of either subspecies degraded the gamma chain.

3.5. Coagulation assay

The serine protease thrombin plays an important role in the blood coagulation cascade by converting native brinogen into brin (Tanaka et al., 2009), a critical step in the polymerization reaction for clot formation. Results of the clot prolongation assay indicate that there was no difference between C. lepidus and C. willardi species ($F_{1,20}$ ¼ 0.147, $p > 0.05$); however, there was a signi cant difference at the subspecies level ($F_{8,20}$ ¼ 34.373, $p < 0.01$). For C. lepidussubspecies, C. l. maculosushad the highest average prolonged clotting time (46 s), followed by C. l. morulus (34 s) and C. l. klauberi (30 s), while C. l. lepidus had the lowest

SF

evaluated in their study. Interestingly, although all C. willardi

Rattlesnake venoms have been classi ed into type I venoms that

Barker, D.G., 1992. Variation, infraspeci c relationships and biogeography of the ridgenose rattlesnake, Crotalus willardi. In: Campbell, J.A., Brodie Jr., E.D. (Eds.), Biology of the Pitvipers. Selva Publishing, Tyler, Texas, pp. 89 e 105.
Barker, D.G., 2016. Ridgenose Rattlesnake, Crotalus willardi (Meek 1906). In: Schuett, G.W., Feldner, M.J., Smith, C.F., Reiserer, R.S. (Eds.), Rattlesnakes of Arizona, vol. 1. ECO Herpetological Publishing, Rodeo, New M exico, pp. 655 e 699.
Blair, C., Sanchez-Ramirez, S., 2016. Diversity-dependent cladogenesis throughout western M exico: evolutionary biogeography of rattlesnakes (Viperidae: Crotalinae: Crotalus and Sistrurus). Mol. Phylogenet. Evol. 97, 145 e 154.
Bryson, R.W., Murphy, R.W., Lathrop, A., Lazcano-Villareal, D., 2011a. Evolutionary drivers of phylogeographical diversity in the highlands of M exico: a case study of the Crotalus triseriatus species group of montane rattlesnakes. J. Biogeogr. 38, 697 e 710.
Bryson, R.W., Murphy, R.W., Graham, M.R., Lathrop, A., Lazcano, D., 2011b. Ephemeral Pleistocene woodlands connect the dots for highland rattlesnakes of the Crotalus intermedius group. J. Biogeogr. 38, 2299 e 2310.
Bryson Jr., R.W., Linkem, C.W., Dorcas, M.E., Lathrop, A., Jones, J.M., Alvarado-Diaz, J., Grünwald, C.I., Murphy, R.W., 2014. Multilocus species delimitation in the Crotalus triseriatus species group (Serpentes: Viperidae: Crotalinae), with the description of two new species. Zootaxa 3826, 475 e 496.
Bryson, R.W., Linkem, C.W., Pavon-Vazquez, C.V., Nieto-Montes de Oca, A., Klicka, J., McCormack, J.E., 2017. A phylogenomic perspective on the biogeography of skinks in the Plestiodon brevirostris group inferred from target enrichment of ultraconserved elements. J. Biogeogr. http://dx.doi.org/10.1111/jbi.12989 .
Calvete, J.J., Sanz, L., Cid, P., De La Torre, P., Flores-Díaz, M., Dos Santos, M.C., Borges, A., Bremo, A., Angulo, Y., Lomonte, B., Alape-Gir on, A., Gutierrez, J.M., 2009. Snake venomics of the Central American rattlesnake Crotalus simus and the South American Crotalus durissus complex points to neurotoxicity as an adaptive paedomorphic trend along Crotalus dispersal in South America. J. Proteome Res. 9, 528 e 544.
Campbell, J.A., Lamar, W.W., 1989. The Venomous Reptiles of Latin America. Cornell University Press, Ithaca, New York .
Campbell, J.A., Lamar, W.W., 2004. The Venomous Reptiles of the Western Hemisphere. Cornell University Press, Ithaca, New York .
Campbell, J.A., 1999. Distribution patterns of amphibians in Middle America. In: Duellman, W.E. (Ed.), Patterns of Distribution of Amphibians: a Global Perspective. Johns Hopkins University Press, Baltimore, pp. 111 e 210.
Castoe, T.A., Sasa, M.M., Parkinson, C.L., 2005. Modeling nucleotide evolution at the mesoscale: the phylogeny of the Neotropical pitvipers of the Porthidium group (Viperidae: Crotalinae). Mol. Phylogenet. Evol. 37, 881 e 898.
Castoe, T.A., Daza, J.M., Smith, E.N., Sasa, M.M., Kuch, U., Campbell, J.A., Chippindale, P.T., Parkinson, C.L., 2009. Comparative phylogeography of pitvipers suggests a consensus of ancient Middle American highland biogeography. J. Biogeogr. 36, 88 e 103.
Dhananjaya, B.L., D'souza, C.J., 2010. An overview on nucleases (DNase, RNase, and phosphodiesterase) in snake venoms. Biochem. Mosc. 75, 1 e 6.
Ferrusquía-Villafranca, I., 1990. Provincias Bi oticas con enfasis en criterios morfotectonicos. Regionalizacion Biogeogra ca, IV.8.10. Atlas Nacional de Mexico, vol. II. Escala 1:4,000,000. Instituto de Geografía, UNAM, M

Noutsos, C., Borevitz, J.O., Hodges, S.A., 2014. Geneow between nascent species: geographic, genotypic and phenotypic differentiation within and between Aquilegia formosa and A. pubescens. Mol. Ecol. 23, 5589–5598.
Place, A.J., Abramson, C.I., 2004. A quantitative analysis of the ancestral area of rattlesnakes. J. Herp. 38, 152–156.
Rael, E.D., Johnson, J.D., Molina, O., McCrystal, H.K., 1992. Distribution of a Mojave toxin-like protein in rock rattlesnake (Crotalus lepidus) venom. In: Campbell, J.A., Brodie Jr., E.D. (Eds.), Biology of the Pitvipers. Selva Publishing, Tyler, Texas, pp. 163–168.
Ramamoorthy, T., Bye, R., Lot, A., Fa, J., 1993. Biological Diversity of Mexico: Origins and Distribution. Oxford University Press, Oxford .
Reed, L.J., Muench, H., 1938. A simple method of estimating fty percent endpoints. Amer. J. Hyg. 27, 493–497.
Rodrigues, R.S., da Silva, J.F., França, J.B., Fonseca, F.P., Otaviano, A.R., Silva, F.H., Hamaguchi, A., Magro, A.J., Braz, A.S.K., dos Santos, J.I., Homsi-Brandeburgo, M.I., Fontes, M.R.M., Fully, A.L., Soares, A.M., Rodrigues, V.M., 2009. Structural and functional properties of Bp-LAAO, a new L-amino acid oxidase isolated from Bothrops pauloensis snake venom. Biochimie 91, 490–501.
Russell, F.E., Beuss, F.W., Woo, M.Y., 1963. Zootoxicological properties of venom phosphodiesterase. Toxicon 1, 99–108.
Samel, M., Vija, H., Rönnholm, G., Siigur, J., Kalkkinen, N., Siigur, E., 2006. Isolation and characterization of an apoptotic and platelet aggregation inhibiting L-amino acid oxidase from Vipera berus berus (common viper) venom. BBA Proteins Proteom. 1764, 707–714.
Saviola, A.J., Pla, D., Sanz, L., Castoe, T.A., Calvete, J.J., Mackessy, S.P., 2015. Comparative venomics of the Prairie Rattlesnake (Crotalus viridis viridis) from Colorado: identi cation of a novel pattern of ontogenetic changes in venom composition and assessment of the immunoreactivity of the commercial anti-venom CroFab®. J. Proteom. 121, 28–43.
Scott Jr., N.J., Reynolds, R.P., 1984. Phenotypic Variation of the Mexican Duck (Anas platyrhynchos Diazi) in Mexico. Condor, pp. 266–274.
Smith, C.F., Mackessy, S.P., 2016. The effects of hybridization on divergent venom phenotypes: characterization of venom from Crotalus scutulatus scutulatus Crotalus oreganus helleri hybrids. Toxicon 120, 110–123.
Souza, D.H., Eugenio, L.M., Fletcher, J.E., Jiang, M.S., Garratt, R.C., Oliva, G., Selistre-de-Araujo, H.S., 1999. Isolation and structural characterization of a cytotoxic L-amino acid oxidase from Agkistrodon contortrix laticinctus snake venom: preliminary crystallographic data. Arch. Biochem. Biophys. 368, 285–290.
Swenson, S., Markland, F.S., 2005. Snake venom brin(ogen)olytic enzymes. Toxicon 45, 1021–1039.
Tanaka, K.A., Key, N.S., Levy, J.H., 2009. Blood coagulation: hemostasis and thrombin regulation. Anesth. Analg. 108, 1433–1446.